

*Michigan*  
**Adult Treatment Protocols**  
 GENERAL PRE-HOSPITAL CARE

Date: November 15, 2012

Page 1 of 1

***General Pre-Hospital Care***

In most cases, the stabilization of patients presenting with medical conditions should be carried out at the patient's side prior to patient movement or transport. Before attempting the following procedures, implement appropriate blood borne and/or airborne pathogen protective procedures. Contact medical control according to local protocol.

Unless otherwise stated, pediatric protocols will apply to patients less than or equal to 14 years of age. If the patient's age is not known, then pediatric protocols will apply until there are physical signs that the patient has reached puberty as indicated by armpit hair in boys and breast development in girls.

**Pre-Medical Control**

**MFR/EMT/SPECIALIST/PARAMEDIC**

1. Assure ABCs while maintaining C-Spine precautions where indicated.
2. Do airway intervention using appropriate airway adjuncts when necessary:

	<b>MFR</b>	<b>EMT</b>	<b>SPECIALIST</b>	<b>PARAMEDIC</b>
<b>Oralpharyngeal Airway</b>	<b>X</b>	<b>X</b>	<b>X</b>	<b>X</b>
<b>Nasopharyngeal Airway</b>	<b>X</b>	<b>X</b>	<b>X</b>	<b>X</b>
<b>Bag-Valve-Mask Ventilation</b>	<b>X</b>	<b>X</b>	<b>X</b>	<b>X</b>
<b>Supraglottic Airway (per MCA approval)</b>		<b>X</b>	<b>X</b>	<b>X</b>
<b>Oral / Nasal Endotracheal Intubation</b>				<b>X/O</b>
<b>Needle / Surgical Cricothyroidotomy</b>				<b>O/O</b>
<b>X: Approved Intervention</b>				
<b>O: Optional Intervention per MCA selection</b>				

3. Administer oxygen and assist ventilations. As indicated refer to the **Emergency Airway Procedure**. Use 2-person BVM technique whenever possible.
4. Obtain an appropriate history and physical exam.
5. Obtain vital signs including pulse oximetry if available or required, approximately every 15 minutes, or more frequently as necessary to monitor the patient's condition (minimum 2 sets suggested).

**SPECIALIST/PARAMEDIC**

6. For pediatric with life threatening or potentially life threatening conditions measure with Broselow Pediatric Emergency Care tape to determine color.
7. Follow specific protocol for patient condition.
8. Establish vascular access per **Vascular Access & IV Fluid Therapy Procedure** when fluid or medication administration may be necessary.

**PARAMEDIC**

9. Apply cardiac monitor and treat rhythm according to appropriate protocol. If available and applicable, obtain 12-lead ECG. A copy of the rhythm strip or 12-lead ECG should be attached to the patient care record and should be left at the receiving facility.
10. Consider use of capnography as appropriate and if available, per **Waveform Capnography Procedure**.

**NOTE:** When possible, take the patient's medications to the hospital.